

Haptics/graphics-facilitated learning and neural recovery

James Patton



Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE DEC 2008		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Haptics/graphics-facilitated learning and neural recovery				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Illinois Chicago				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002187. Proceedings of the Army Science Conference (26th) Held in Orlando, Florida on 1-4 December 2008, The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 35	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Research at the RIC




Can you use robotics and/or display feedback technology to train movement skills?




Haptic/Graphic
Interaction
with
Simple Objects

Detecting Hemispatial Neglect in stroke survivors (Assaf Dvorkin)



**HEALTHY
CONTROL**



**PATIENT WITH
NEGLECT**



Correct detection



Interactive technology can

N o v e l t y

- Give Precision guidance
- Assist as needed; wean
- Track & store progress
- Rapidly present scenarios
- Render “unreal” forces
- Challenge (train robustness)
- Distort reality & cause adaptation
- Possibly be worn and/or taken home


Long latencies make feedback control impossible for most everyday movements

<i>REFLEX TYPE:</i>	<i>Movement Latencies</i>	<i>Frequencies</i>
<i>Musculoskeletal Impedances</i>	<i>instantaneous</i>	∞
<i>Spinal</i>	<i>30-80 msec (Dewhurst, 1967)</i>	<i>1.7 Hz (Hogan, 1990)</i>
<i>Triggered</i>	<i>80-120 msec (Crago, et al., 1976)</i>	<i>0.6 Hz (Hogan, 1990)</i>
<i>Supraspinal (long)</i>	<i>120-180 msec (Schmidt, 1988)</i>	<i>0.5 Hz (Hogan, 1990)</i>
<i>Vision</i>	<i>100 msec (Nashner and Berthoz, 1978)</i>	<i>0.6 Hz (Hogan, 1990)</i>
<i>Vestibular</i>	<i>102 msec (Melvill Jones and Watt, 1971)</i>	<i>0.6 Hz (Hogan, 1990)</i>


Background on robotic force field training

adaptation and after-effects

PROGRESSIVE TRAINING:



THEN TURN OFF THE FORCES:



"After-effects"

*Shadmehr, R and Mussa-Ivaldi, FA (1994)
Journal of Neuroscience 14: 3208-3224.*

Dynamic Model of the Arm & controller

Functional form assumptions

$$\underbrace{I(x)\ddot{x} + G(x)}_D = 0 \quad (\text{uncontrolled})$$

$$D - C = 0 \quad (\text{controlled})$$

$$\underbrace{\{I(x)\ddot{x} + G(x, \dot{x})\}}_D - \underbrace{\left\{ \underbrace{\hat{I}(x_{E(t)})\ddot{x}_{E(t)} + \hat{G}(x_{E(t)}, \dot{x}_{E(t)})}_{C_{FF}} + \underbrace{Z[x_{E(t)} - x]}_{C_Z} \right\}}_C = 0$$











$$\underbrace{\hspace{10em}}_{M(x, \dot{x}, \ddot{x}, x_{E(t)} \mid 18 \text{params})}$$


simplest learning rule:

$$\tau_i = \tau_{i-1} + \mu(e_{i-1})$$

Adaptive Training

Techniques for possibly facilitating learning

-  • **Human-human collaboration**
-  • **Custom-designed force fields**
-  • **Custom-designed Visual distortions**
-  • **Error augmentation (force and vision)**
-  • **Obstacle avoidance (changes desired traj)**
-  • **Sensory crossover-teach visual w/force**
-  • **Gradual learning**
-  • **Stochastic Resonance (injected noise)**
-  • **Intermanual and bimanual transfer**
-  • **Manipulation of stability limits**



Early
exposure





**ALTERING
MOVEMENTS
IN THE
HEALTHY**


Late
Training

After-
Effects

Late
Washout


Mussa-Ivaldi, F.A., Patton, J.
L., IEEE International
Conference on Robotics and
Automation, 2000, San
Francisco, CA






Patton, J. L., M. E. Phillips-Stoykov, et al. (2006). "Evaluation of robotic training forces that either enhance or reduce error in chronic hemiparetic stroke survivors." *Experimental Brain Research* 168(3): 368-383.

Performance Improvement vs Enhancement of error




Patton, J. L., M. E. Phillips-Stoykov, et al. (2006). "Evaluation of robotic training forces that either enhance or reduce error in chronic hemiparetic stroke survivors." *Experimental Brain Research* 168(3): 368-383.


A. Unperturbed baseline




B. Late Machine learning




C. Early training




D. Late training




E. after-effects




F. early washout



G. late washout




(Subject sfd54)



Custom-Designed Training Forces:

Error reduces significantly in stroke survivors


Patton Et Al (2005) Custom-designed haptic training for restoring reaching ability to individuals with stroke. *JRRD* in press




Error Augmentation


ERROR AUGMENTATION CANDIDATES

GAIN





OFFSET






Error Augmentation speeds up & increases learning in healthy individuals




Current Work: 2 hands

- Therapist-Driven Trajectories
- Patient-Driven Trajectories




Free exploration learning with error augmentation via negative damping enhances learning


(Felix Huang)





A. Planar manipulandum presents anisotropic inertial



B. Interactive Priming Phase




C. Performance Evaluation Phase




Applications for adaptive training

- **Rehabilitation**
- **Teleoperation**
- **Human-machine interactions**
- **Learning and Co-Learning a Neural Machine interfaces**






Thanks

- **W. Z. Rymer**
- **F. A. Mussa-Ivaldi**
- **F. Huang**
- **A. Dvorkin**
- **R. Kenyon**
- **Y. Wei**
- **M. Peshkin**
- **M. E. Phillips**
- **M. Kovic**
- **R. Haner**
- **C. Malecka**
- **P. Shah**
- **C. Raasch**
- **L. Kahn**
- **D. Sha**
- **NIH R01 R01 NS053606**
- **NIDRR H133A080045**
- **NIDRR H133E0700 13**
- **AHA 0330411Z**
- **NIH R24 HD39627**
- **NIH 5 T32 HD07418**
- **NIH 5 RO1 NS 35673**
- **NIH F32HD08658**
- **NIDRR RERC 0330411Z**
- **Falk Trust**
- **Davee Foundation**




Sensory Crossover







Initial direction error with desired for movement direction of 315degrees



Assessments

- Blinded Rater
- Assessments pre and post each tx.
 - “Reach and retrieve” (rag on a stick)
 - Functional workspace when reaching towards 9 targets on periphery
 - Wolf Motor Function Test
 - Box and Blocks
 - Fugl-Meyer




Figure 3: Schematic example of the directions and layers of the functional workspace determined in the workspace parts of the experiment.

Secondary assessments:

- How much time (tx vs. setup)?
- how long did it take to achieve therapeutically meaningful effect?
- Which treatment engaged / frustrated patient
- Therapist opinion

EA haptic forces




Figure 11: The double exponential function governing the error-augmentation (dashed line defined in Equation 1).

Inclusion Criteria

- Chronic Stroke (8+ mos post) (*Changed to 6+ mos in NW IRB*)
- Available medical records/radiographic info
- Ischemic infarct in MCA (*not included in NW IRB*)
- Primary Motor Cortex Involvement (*not included in NW IRB*)
- Some degree of shoulder and elbow mvmt.
- AMFM 40-50 (*Changed to 25-50 in NW IRB*)

Exclusion Criteria

- Bilateral paresis
- Severe sensory deficits
- Severe spasticity (MAS = 4)
- Severe contracture (*added to NW IRB*)
- Aphasia, cognitive impairment or affective dysfunction that would influence the ability to perform experiment
- Severe concurrent medical problems
- Diffuse/multiple lesion sites or multiple stroke events
- Hemispatial neglect/inattention or field cut that would influence the ability to perform experiment
- Ataxia (*added in NW IRB*)
- Significant pain (greater than 5/10) in UE (*added in NW IRB*)
- Botox injection in previous 3 months (*added in NW IRB*)
- Participation in other UE research projects (*added in NW IRB*)

Statistics

- Randomized Mixed effects model
 - Trend affecting the hypothesis on treatment type?
 - Period-by treatment interaction
 - Carry over effect
 - Patient-by-treatment interaction
- Bayesian?
 - Early results
 - Alternative to testing

R1: AIM 1, Experiment 1.1

- Therapist-Driven Trajectories
- 17 subjects, random order of groups
- 4 weeks of tx, 1 treatment per week
- (45 min)
- 4 groups
 - EA with visual distortion
 - EA with haptic forces
 - EA with visual distortion and haptics
 - No EA




Figure 11: The double exponential function governing the error-augmentation (dashed line defined in Equation 1).